arXiv:hep-th/9207090v1 28 Jul 1992

INVARIANTS OF COLORED LINKS AND A PROPERTY OF THE CLEBSCH-GORDAN COEFFICIENTS OF $U_q(g)$[1]

TETSUO DEGUCHI and TOMOTADA OHTSUKI †

Department of Physics, Faculty of Science,
University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113, Japan

† *Department of Mathematical Sciences,*
University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113, Japan

Abstract.

We show that multivariable colored link invariants are derived from the roots of unity representations of $U_q(g)$. We propose a property of the Clebsch-Gordan coefficients of $U_q(g)$, which is important for defining the invariants of colored links. For $U_q(sl_2)$ we explicitly prove the property, and then construct invariants of colored links and colored ribbon graphs, which generalize the multivariable Alexander polynomial.

1 Introduction

Recently a new family of invariants of colored oriented links and colored oriented ribbon graphs is introduced, which gives generalizations of the multivariable Alexander polynomial.[1,2,6] The new invariants are related to the roots of unity representations of $U_q(sl_2)$ where $(X^{\pm})^N = 0$ and $(K)^N = q^{2Np}$ ($p \in \mathbf{C}$) and $q = \epsilon$.[5,6,7] Here $\epsilon = \exp(\pi i s/N)$, and the integers N and s are coprime. We call the representations nilpotent representations.

The invariants of colored links have a property that they vanish for disconnected links.[2,6] Due to this property a proper regularization method is necessary for definition of the invariants.

In this paper we show an important property of the Clebsch-Gordan coefficients (CGC) of the nilpotent representations of $U_q(sl_2)$, which leads to the definition of the colored link invariants. We consider the nilpotent reps of $U_q(g)$ such that $(X_i^{\pm})^N = 0$, $(K_i)^N = q^{2Np_i}$ ($p_i \in \mathbf{C}$) and $q = \epsilon$. We give a conjecture that the property of CGC also holds for the nilpotent reps of $U_q(g)$ and we can define invariants of colored links for $U_q(g)$ in the same way as $U_q(sl_2)$.

2 CGC of the nilpotent representations

We introduce some symbols for a positive integer n and a complex parameter p.

$$[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}, \quad [n]_q! = \prod_{k=1}^{n} [k]_q, \quad [p;n]_q! = \prod_{k=0}^{n-1} [p-k]_q. \tag{1}$$

---

[1] to appear in the Proceedings of the 21st International Conference on the Differential Geometry Methods in Theoretical Physics, 5-9 June, 1992, Tianjin, China

For $n = 0$ we assume $[0]_q! = [p;0]_q! = 1$.

We introduce the nilpotent rep. $V^{(p)}$ [5]: $\pi(X^+)^a_b = \sqrt{[2p-a]_\epsilon [a+1]_\epsilon}\ \delta_{a+1,b}$, $\pi(X^-)^a_b = \sqrt{[2p-a+1]_\epsilon [a]_\epsilon}\ \delta_{a-1,b}$, $\pi(K)^a_b = \epsilon^{(2p-2a)}\delta_{a,b}$. Here $a, b = 0, 1, \cdots N-1$. We note that N is related to the dimensions of the representation. When N is even, ϵ is $2N$-th primitive root of unity, while when N is odd, ϵ may be $2N$-th or N-th primitive root of unity.

In Ref. 1 the explicit matrix representations of the colored braid group were introduced. It was shown that the representations are equivalent to the R matrices of the nilpotent reps, which are derived from the universal R matrix $\mathcal{R}$: $R^{a_1,a_2}_{b_1,b_2} = \pi^{p1} \otimes \pi^{p_2}(\mathcal{R})^{a_1 a_2}_{b_1 b_2}$. [5]

We can show the fusion rule for the tensor product: $V^{(p_1)} \otimes V^{(p_2)} = \sum_{p_3} N^{p_3}_{p_1 p_2} V^{(p_3)}$, where $N^{p_3}_{p_1,p_2} = 1$ for $p_3 = p_1 + p_2 - n$ ($0 \leq n \leq N-1$ and $n \in \mathbf{Z}$); $N^{p_3}_{p_1,p_2} = 0$, otherwise. The CGC for $V^{(p_i)}$ $(i = 1, 2, 3)$ are given by [6]

$$\begin{aligned}
&C(p_1, p_2, p_3; z_1, z_2, z_3) = \delta(z_3, z_1 + z_2 - n) \\
&\times \sqrt{[2p_1 + 2p_2 - 2n + 1]_\epsilon} \sqrt{[n]_\epsilon! [z_1]_\epsilon! [z_2]_\epsilon! [z_3]_\epsilon!} \\
&\times \sum_\nu \frac{(-1)^\nu \epsilon^{-\nu(p_1+p_2+p_3+1)} \times \epsilon^{(n-n^2)/2+(n-z_2)p_1+(n+z_1)p_2}}{[\nu]_\epsilon! [n-\nu]_\epsilon! [z_1-\nu]_\epsilon! [z_2-n+\nu]_\epsilon!} \\
&\times \sqrt{\frac{[2p_1-n; z_1-\nu]_\epsilon! [2p_1-z_1; n-\nu]_\epsilon! [2p_2-n; z_2+\nu-n]_\epsilon! [2p_2-z_2; \nu]_\epsilon!}{[2p_1+2p_2-n+1; z_1+z_2+1]_\epsilon!}}. \quad (2)
\end{aligned}$$

Here $0 \leq z_i \leq N-1$, for i=1,2,3, and the sum over the integer ν in (2) is taken under the condition: max $\{0, n-z_2\} \leq \nu \leq$ min $\{n, z_1\}$. The expression of the CGC was proved through the infinite dimensional representations. [6]

3 A Property of CGC

We consider the following two sums;

$$\begin{aligned}
A^n_{p_1 p_2} &= \sum_{z_1} \left[C(p_1, p_2, p_3; z_1, z_2, z_1 + z_2 - n)\right]^2 q^{2\rho(z_1)}, \\
B^n_{p_1 p_2} &= \sum_{z_2} \left[C(p_1, p_2, p_3; z_1, z_2, z_1 + z_2 - n)\right]^2 q^{-2\rho(z_2)}. \quad (3)
\end{aligned}$$

For the nilpotent reps of $U_q(sl_2)$ we assume $\rho(z) = p - z + Np/2$ and $q = \epsilon$. Due to the irreducibility of the representations the sum $A^n_{p_1 p_2}$ $(B^n_{p_1 p_2})$ does not depend on z_2 (z_1). We now introduce an important property of CGC.

Proposition 3.1 *The CGC of the nilpotent representations satisfy*

$$A^n_{p_1 p_2}/B^n_{p_1 p_2} = g(p_1, p_2) = f(p_1)/f(p_2), \ \textit{independent of } n \quad (p_3 = p_1 + p_2 - n). \quad (4)$$

(For the nilpotent reps we have $g(p_1, p_2) = [2p_2; N-1]_\epsilon! / [2p_1; N-1]_\epsilon!$).

Proof

We set $z_2 = 0$ for $A^n_{p_1 p_2}$ and $z_1 = 0$ for $B^n_{p_1 p_2}$, and we define $L^n_{p_1 p_2}$ and $R^n_{p_1 p_2}$ by $A^n_{p_1 p_2} = L^n_{p_1 p_2}[2p_3+1]/[2p_3+1+n; N]!$ and $B^n_{p_1 p_2} = R^n_{p_1 p_2}[2p_3+1]/[2p_3+1+n; N]!$, respectively. Then we can show

$$L^n_{p_1 p_2} = (-1)^{nN}[2p_2; N-1]_\epsilon!, \quad R^n_{p_1 p_2} = (-1)^{nN}[2p_1; N-1]_\epsilon!. \tag{5}$$

The first eq. in (5) can be shown by using the following recurrence relation on n.

$$L^n_{p_1 p_2} = \frac{[2p_1 + 2p_2 - n - N + 2]}{[2p_2+1]} \epsilon^{2p_1 - n} L^{n+1}_{p_1+1/2, p_2+1/2} - \frac{[2p_1 - n]}{[2p_2+1]} L^{n+1}_{p_1 p_2 + 1/2}. \tag{6}$$

The second eq. in (5) (for $R^n_{p_1 p_2}$) is derived from that of $L^n_{p_1 p_2}$ by exchanging p_1 with p_2, and by setting $\epsilon \to \epsilon^{-1}$.

It is easy to see that the property (4) of CGC also holds for the finite dimensional (spin) representations of $U_q(sl_2)$ with q generic, where we have $g(j_1, j_2) = [2j_1]/[2j_2]$.

We can define invariants of colored links using the proposition 3.1. Let T be a $(1,1)$- tangle. We denote by $\hat{T}$ the link obtained by closing the open strings of T. We note the following proposition. [2]

Proposition 3.2 *Let T_1 and T_2 be two $(1,1)$-tangles. If $\hat{T}_1$ is isotopic to $\hat{T}_2$ as a link in S^3 by an isotopy which carries the closing component of $\hat{T}_1$ to that of $\hat{T}_2$. Then T_1 is isotopic to T_2 as a $(1,1)$-tangle.*

Let us introduce the functor $\phi(\cdot)$ for the tangle diagrams. [2] We denote by $\phi(T, \boldsymbol{\alpha})^a_b$ the value ϕ for the tangle with variables a and b on the closing component (or edge). It is easy to show that $\phi(T, \boldsymbol{\alpha})^a_b = \lambda \delta_{ab}$. [2]

We put $L = \hat{T}$ and s is the color of the closing component (or edge) of $\hat{T}$. For a colored link $(L, \boldsymbol{\alpha})$ and a color s of closing component (or edge), we define Φ by $\Phi(L, s, \boldsymbol{\alpha}) = \lambda$ where L, T, s are as above and $\phi(T, \boldsymbol{\alpha})^a_b = \lambda \delta_{ab}$. We can show that Φ is well-defined, i.e. $\Phi(L, s, \boldsymbol{\alpha})$ does not depend on a choice of T. [2] From the proposition 3.1 we have the following.

Proposition 3.3 [2] *For a link L and its color $\boldsymbol{\alpha} = (p_1, \cdots, p_n)$, we have*

$$\Phi(L, s, \boldsymbol{\alpha})([p_s; N-1]_\epsilon!)^{-1} = \Phi(L, s', \boldsymbol{\alpha})([p_{s'}; N-1]_\epsilon!)^{-1}. \tag{7}$$

Definition 3.4 [2] *For a colored oriented link $(L, \boldsymbol{\alpha})$, we define an isotopy invariant $\hat{\Phi}$ of $(F, \boldsymbol{\alpha})$ by*

$$\hat{\Phi}(L, \boldsymbol{\alpha}) = \Phi(L, s, \boldsymbol{\alpha})([p_s; N-1]_\epsilon!)^{-1}. \tag{8}$$

Thus we have constructed the multivariable invariants from the property (4) of CGC. In the same way we can define the multivariable invariants of colored ribbon graphs. [6]

We now consider CGC of $U_q(g)$, where g is a simple Lie algebra.

$$\begin{aligned} A^{\Lambda_3}_{\Lambda_1\Lambda_2} &= \sum_{\vec{z}_1} \left[C(\Lambda_1,\Lambda_2,\Lambda_n;\vec{z}_1,\vec{z}_2,\vec{z}_1+\vec{z}_2-\vec{n})\right]^2 q^{2\rho(\vec{z}_1)}, \\ B^{\Lambda_3}_{\Lambda_1\Lambda_2} &= \sum_{\vec{z}_2} \left[C(\Lambda_1,\Lambda_2,\Lambda_3;\vec{z}_1,\vec{z}_2,\vec{z}_1+\vec{z}_2-\vec{n})\right]^2 q^{-2\rho(\vec{z}_2)}. \end{aligned} \tag{9}$$

We assume that ρ is given by "half the sum of positive roots". Finally, we propose the following conjecture.

Conjecture 3.5 *(1) The CGC of the nilpotent representations Λ_i of $U_q(g)$ satisfy*

$$A^{\Lambda_3}_{\Lambda_1\Lambda_2}/B^{\Lambda_3}_{\Lambda_1\Lambda_2} = g(\Lambda_1,\Lambda_2), \text{ independent of } \Lambda_3. \tag{10}$$

(2) With a proper normalization of the quantum trace ($\rho(\vec{z}) \to \rho(\vec{z})$ + constant), we can set $g(\Lambda_1,\Lambda_2) = f(\Lambda_1)/f(\Lambda_2)$.
(3) $f(\Lambda)$ is equivalent to the tangle invariant for the Hopf link.

If the conjecture is true, we can construct multivariable invariants of colored links and colored ribbon graphs from $U_q(g)$ in the same way as $U_q(sl_2)$.

It is easy to show that the property (10) holds for CGC of finite dimensional representations of $U_q(g)$ with q generic.

References

[1] Y. Akutsu and T. Deguchi, A New Hierarchy of Colored Braid Group Representations, Phys. Rev. Lett. **67** (1991) 777.

[2] Y. Akutsu, T. Deguchi and T. Ohtsuki, Invariants of Colored Links, Journal of Knot Theory and Its Ramifications **1** (1992) pp. 161-184.

[3] T. Deguchi and Y. Akutsu, A General Formula for Colored Z_N Graded Braid Matrices and Fusion Braid Matrices, J. Phys. Soc. Jpn. **60** (1991) 2559.

[4] T. Deguchi and Y. Akutsu, A New Hierarchy of Colored Vertex Models, J. Phys. Soc. Jpn. **60** (1991) 4053.

[5] T. Deguchi and Y. Akutsu, Colored braid matrices from infinite dimensional representations of $U_q(g)$, Mod. Phys. Lett. **A7** (1992) 767.

[6] T. Deguchi and Y. Akutsu, Colored Vertex Models, Colored IRF Models, and Invariants of Colored Framed Graphs, preprint January 1992.

[7] T. Ohtsuki, Colored ribbon Hopf algebras and universal invariants of framed links, preprint June 1992.